

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68098 A2

(51) International Patent Classification⁷: **A61K 31/55**,
A61P 35/00

(74) Agents: **TRECARTIN, Richard, F. et al.**; Flehr Hohbach
Test Albritton & Herbert LLP, 4 Embarcadero Center, Suite
3400, San Francisco, CA 94111-4187 (US).

(21) International Application Number: **PCT/US01/08480**

(22) International Filing Date: **15 March 2001 (15.03.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/189,699 **15 March 2000 (15.03.2000) US**

(71) Applicant (*for all designated States except US*): **CHEM-
GENEX THERAPEUTICS, INC.** [US/US]; 34700 Cam-
pus Drive, Fremont, CA 94555 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BROWN, Dennis,**
M. [US/US]; 100 San Mateo Drive, Menlo Park, CA 94025
(US). **MICHAELS, Shawnya, J.** [US/US]; 206 Dolores
Street, El Granada, CA 94018 (US).

(54) Title: **CEPHALOTAXINE ALKALOID COMPOSITIONS AND USES THEREOF**

(57) Abstract: A method of treatment of a host with a cellular proliferative disease, comprising contacting the host with a cephalotax-
ine and an antiproliferative agent, each in an amount sufficient to modulate said cellular proliferative disease, is described. In some
embodiments, the cephalotaxine comprises homoharringtonine (cephalotaxine, 4-methyl-2-hydroxy-2-(4-hydroxy-4-methyl pentyl)
butanedioclate ester). Antiproliferative agents of the invention comprise alkylating agents, intercalating agents, metal coordination
complexes, pyrimidine nucleosides, purine nucleosides, inhibitors of nucleic acid associated enzymes and proteins, and agents af-
fecting structural proteins and cytoplasmic enzymes.

WO 01/68098 A2

CEPHALOTAXINE ALKALOID COMPOSITIONS AND USES THEREOF

5 This application claims the benefit of U.S. Provisional Application Number 60/189,699, filed March 15, 2000.

FIELD OF THE INVENTION

The technical field of the invention is the use of cephalotaxine alkaloids with antiproliferative agents to treat a host with a cellular proliferative disease.

BACKGROUND OF THE INVENTION

10 There is considerable interest in modulating the efficacy of currently used antiproliferative agents to increase the rates and duration of antitumor effects associated with conventional antineoplastic agents.

15 Conventional antiproliferative agents used in the treatment of cancer are broadly grouped as chemical compounds which (1) affect the integrity of nucleic acid polymers by binding, alkylating, inducing strand breaks, intercalating between base pairs or affecting enzymes which maintain the integrity and function of DNA and RNA; (2) chemical agents that bind to proteins to inhibit enzymatic action (*e.g.*, antimetabolites) or the function of structural proteins necessary for cellular integrity (*e.g.*, antitubulin agents). Other chemical compounds
20 that have been identified to be useful in the treatment of some cancers include drugs which block steroid hormone action for the treatment of breast and prostate cancer, photochemically activated agents, radiation sensitizers and protectors.

Of special interest to this invention are those compounds that directly affect the integrity of

the genetic structure of the cancer cells. Nucleic acid polymers such as DNA and RNA are prime targets for anticancer drugs. Alkylating agents such as nitrogen mustards, nitrosoureas, aziridine containing compounds directly attack DNA. Metal coordination compounds such as cisplatin and carboplatin similarly directly attack the nucleic acid structure resulting in lesions that are difficult for the cells to repair which, in turn, can result in cell death. Other nucleic acid affecting compounds include anthracycline molecules such as doxorubicin, which intercalates between the nucleic acid base pairs of DNA polymers, bleomycin which causes nucleic acid strand breaks, and fraudulent nucleosides. Fraudulent nucleosides include pyrimidine and purine nucleoside analogs which are inappropriately incorporated into nucleic polymer structures and ultimately cause premature DNA chain termination. Certain enzymes that affect the integrity and functionality of the genome can also be inhibited in cancer cells by specific chemical agents and result in cancer cell death. These include enzymes that affect ribonucleotide reductase (*e.g.*, hydroxyurea, gemcitabine), topoisomerase I (*e.g.* camptothecin) and topoisomerase II (*e.g.*, etoposide).

One of the most broadly used of these DNA targeted anticancer drugs is cisplatin (cis-diamminedichloroplatinum II, CDDP). This compound is active against several human cancers including testicular, small-cell lung, bladder, cervical and head and neck cancer.

Although the clinical activity of currently approved antiproliferative agents against many forms of cancers can be shown, improvements in tumor response rates, duration of response and ultimately patient survival are still sought. The invention described herein demonstrates the novel use of the cephalotaxine alkaloids and analogs thereof, including homoharringtonine (HHT) which can potentiate the antitumor effects of chemotherapeutic drugs, in particular, agents affecting the integrity of nucleic polymers such as DNA.

SUMMARY OF THE INVENTION

Methods and compositions are provided for the treatment of a host with a cellular proliferative disease, particularly a neoplasia. In the subject methods, pharmaceutically

acceptable cephalotaxine and an antiproliferative agent are administered in an amount sufficient to modulate the cellular proliferative disease.

DETAILED DESCRIPTION OF THE FIGURES

Figure 1 depicts the general structure of a cephalotaxine analog. R1 and R2 represent substitution groups. Structures for R1 and R2 are shown for the cephalotaxine analog, Homoharringtonine.

Figure 2 depicts the structure of the cephalotaxine analog, Homoharringtonine.

Figure 3 shows tumor growth delay, as tumor volume on days after treatment with HHT, HHT followed by CDDP, or CDDP alone.

DETAILED DESCRIPTION OF THE INVENTION

Methods and compositions are provided for the treatment of a host with a cellular proliferative disease, particularly a neoplasia. In the subject methods, a pharmaceutically acceptable cephalotaxine is administered, preferably systemically, in conjunction with an antiproliferative agent to improve the anticancer effects. In a preferred embodiment, the cephalotaxine provides a chemopotentiator effect.

The agents are provided in amounts sufficient to modulate a cellular proliferative disease. In one embodiment, modulation of a cellular proliferative disease comprises a reduction in tumor growth. In another embodiment, modulation of a disease comprises inhibition of tumor growth. In another embodiment, modulation of a cellular proliferative disease comprises an increase in tumor volume quadrupling time (described below). In another embodiment, modulation of a cellular proliferative disease comprises a chemopotentiator effect. In another embodiment, modulation of a disease comprises a chemosensitizing effect. In other embodiments, modulation of a disease comprises cytostasis. In still other embodiments, modulation of a disease comprises a cytotoxic effect.

A chemical agent is a "chemopotentiator" when it enhances the effect of a known antiproliferative drug in a more than additive fashion relative to the activity of the chemopotentiator or antiproliferative agent used alone. In some cases, a chemosensitizing effect may be observed. This is defined as the effect of use of an agent that if used alone would not demonstrate significant antitumor effects but would improve the antitumor effects of an antiproliferative agent as compared to use of the antiproliferative agent by itself.

As used herein, the term "cephalotaxine" includes all members of that chemical family including alkaloid derivatives of the Chinese evergreen, *Cephalotaxus fortunei* and analogs thereof. The cephalotaxine family is defined by chemical structure as the ring structures in Figure 1.

A cephalotaxine analog is further defined but not limited to the structure depicted in Figure 1, having substituent or substitute groups at R1 and R2. Examples of R1 and/or R2 include esters, including herringtonine, isoharringtonine, homoharringtonine, deoxyharringtonine, acetylcephalotaxine and the like. Table 1 lists structures of R1 and R2 for some of these analogs. R1 and R2 substitutions are typically employed to improve biological activity, pharmaceutical attributes such as bioavailability or stability, or decrease toxicity. In one embodiment, R1 and/or R2 include alkyl substitutions (e.g., methyl, ethyl, propyl etc.). In another embodiment, R1 and/or R2 include esters (e.g., methoxy, ethoxy, butoxy, etc.). R1 and R2 are not limited to the above examples, however, in the scope of this invention.

Table 1

	R1	R2
5 isoharringtonine	-OCH ₃	$\begin{array}{c} \text{OH} \quad \text{OH} \\ \quad \\ \text{CH}_3\text{CH}(\text{CH}_2)_2\text{C} - \text{CHCO}_2\text{CH}_3 \\ \\ \text{CO}_2^- \end{array}$
10 harringtonine	-OCH ₃	$\begin{array}{c} \text{OH} \quad \text{OH} \\ \quad \\ \text{CH}_3\text{C}(\text{CH}_2)_2\text{C} - \text{CH}_2\text{CO}_2\text{CH}_3 \\ \quad \\ \text{CH}_3 \quad \text{CO}_2^- \end{array}$
acetylcephalotaxine	-OCH ₃	CH ₃ CO ₂ ⁻
15 homoharringtonine	-OCH ₃	$\begin{array}{c} \text{OH} \quad \text{OH} \\ \quad \\ \text{CH}_3\text{C}(\text{CH}_2)_3\text{C} - \text{CH}_2\text{CO}_2\text{CH}_3 \\ \quad \\ \text{CH}_3 \quad \text{CO}_2^- \end{array}$

A cephalotaxine analog is a further chemical refinement. A specific example of a cephalotaxine analog is homoharringtonine which is the butanediocate ester of cephalotaxine, 4-methyl-2-hydroxy-2-(4-hydroxy-4-methyl pentyl) (Figure 2).

As used herein, antiproliferative agents are compounds which induce cytostasis or cytotoxicity. "Cytostasis" is the inhibition of cells from growing while "cytotoxicity" is defined as the killing of cells. Specific examples of antiproliferative agents include: antimetabolites, such as methotrexate, 5-fluorouracil, gemcitabine, cytarabine, pentostatin, 6-mercaptapurine, 6-thioguanine, L-asparaginase, hydroxyurea, N-phosphonoacetyl-L-aspartate (PALA), fludarabine, 2-chlorodeoxyadenosine, and floxuridine; structural protein agents, such as the vinca alkaloids, including vinblastine, vincristine, vindesine, vinorelbine paclitaxel, and colchicine; antibiotics, such as dactinomycin, daunorubicin, doxorubicin,

idarubicin, bleomycins, plicamycin, and mitomycin; hormone antagonists, such as tamoxifen and luteinizing hormone releasing hormone (LHRH) analogs; nucleic acid damaging agents such as the alkylating agents mechlorethamine, cyclophosphamide, ifosfamide, chlorambucil, dacarbazine, methylnitrosourea, semustine (methyl-CCNU), chlorozotocin, busulfan, procarbazine, melphalan, carmustine (BCNU), lomustine (CCNU), and thiotepa, the intercalating agents doxorubicin, dactinomycin, daurorubicin and mitoxantrone, the topoisomerase inhibitors etoposide, camptothecin and teniposide, and the metal coordination complexes cisplatin and carboplatin.

The following examples are offered by way of illustration and not by way of limitation.

EXAMPLES

Example 1

Chemopotential of Cisplatin (CDDP) by Homoharringtonine (HHT)

Transplantable experimental murine fibrosarcomas (2×10^5 RIF-1 cells) were grown intradermally in the flanks of 3 month old female C3H mice (Charles River, Holister, CA).

When the tumors reached a volume of approximately 100 mm^3 , the mice were randomly assigned to each experimental group (4 mice per group).

The experimental compositions were prepared as described in Table 2.

Table 2

Agent	Dose	Solvent	Supplier
Homoharringtonine	2 mg/kg	DMSO	NCI
Cisplatin	4 mg/kg	Water for injection	David Bull Labs

The chemopotentiator, homoharringtonine, was obtained from NCI and was made to the

appropriate concentration in DMSO. Cisplatin (David Bull Laboratories- Mulgrave, Australia, lot. 5201844x) was made to the appropriate concentration in water for injection. The compositions were injected systemically (i.e., intraperitoneally, i.p.), in a volume of 100 microliters. For the treatment of group 3, the chemopotentiator, homoharringtonine, was injected 30 minutes prior to the injection of cisplatin. After treatment, the growth of the tumors was monitored three times per week by caliper measurements of three perpendicular diameters of the tumor and calculation of tumor volume from the formula:

$$V = \pi / 6 \times D_1 \times D_2 \times D_3,$$

where D_{1-3} is in mm.

The tumors were followed until they reached a size of four times their day zero treatment volume (TVQT), or up to 30 days after treatment, whichever came first. The data is expressed as the "tumor volume quadrupling time" (TVQT) mean and as the "delay." Mean TVQT is the mean days required for individual tumors to grow to four times the tumor volume at the initial treatment day. The "delay" is the median of days required for a tumor to grow to four times the mean size of the treated group, minus the median of days required to grow to four times the mean size of the control group. The data is also expressed as the ratio of the tumor volume quadrupling time of the treated tumor over the untreated control group (TVQT/CTVQT). Increasing values of this ratio indicate increased antitumor response.

The data is presented in Table 3 below and in Figure 3.

Table 3

Group	Treatment	Dose (mg/kg)	Mean TVQT ± S.E.	TVQT/CTVQT	Median (TVQT)	Delay (Days)
1	Untreated Control	-	8.3 ± 0.4	1.0	8.6	0.00
2	Homoharringtonine	2	10.1 ± 0.4	1.2	9.8	1.20
3	Homoharringtonine → Cisplatin	2 → 4	14.9 ± 0.8	1.8	14.8	6.17
4	Cisplatin	4	12.9 ± 1.1	1.5	12.5	3.83

The arrow (→) in Group 3 indicates administration 30 minutes following administration of homoharringtonine.

The results of Table 3 indicate that the antiproliferative activity of cisplatin is enhanced by the use of the chemopotentiator, homoharringtonine in that a more than additive effect was observed when both compounds were used to treat the tumor bearing mice (group 3) in comparison to the use of cisplatin alone (group 4) or homoharringtonine alone (group 2).

5 Example 2

Effect of Homoharringtonine, Alone and in Combination with Other Chemotherapeutics on RIF-1 Tumor Growth in C3H Mice

The RIF-1 murine fibrosarcoma tumor model was used to evaluate the antitumor activity of homoharringtonine, alone and in combination with various antiproliferative agents. The
10 antiproliferative agents used include those that affect nucleic acid (*e.g.*, DNA) integrity (*e.g.*, cisplatin, cytarabine, camptothecin, etoposide, 5-fluorouracil, or amonafide), agents that affect structural proteins (*e.g.*, paclitaxel, vinblastine, or colchicine) or cytoplasmic enzymes (*e.g.*, genistein).

Homoharringtonine (HHT-NCI) was obtained from NCI as a powder. Homoharringtonine
15 (HHT-Clin) was obtained from Hangzhou Minsheng Pharmaceutical Group (Hangzhou, China), in 1 mL vials, prediluted with water to 1 mg/mL. Cisplatin for Injection, USP, was obtained from David Bull Labs (Mulgrave, Australia), Lot No.5201844x, as a lyophilized powder. Paclitaxel was obtained from Bristol Myers Squibb Co. (Princeton, NJ), Lot No.
9J16241, exp. Sep 2001, prediluted to 6 mg/mL in Cremaphor/EL. Cytarabine was obtained
20 from Bedford Labs (Bedford, OH), Lot No.86968A, exp. 6/02, as a lyophilized powder. Camptothecin was obtained from Boehringer-Ingelheim, Lot No.142088, as a powder. Vinblastine was obtained from Bedford Labs (Bedford, OH), Lot No.112647, as a lyophilized powder. Etoposide was obtained from Pharmacia (Kalamazoo, MI), Lot No. ETA013, exp.
5/99, as a liquid prediluted to 20 mg/mL. 5-Fluorouracil was obtained from Pharmacia
25 (Kalamazoo, MI), Lot No.FFA191, exp. 7/00, as a liquid prediluted to 50 mg/mL. Amonafide was obtained from Penta Biotech (Union City, CA), Lot No.039-01, as a powder.

Colchicine was obtained from Sigma (St. Louis, MO), Lot No.55H0685, as a powder.
Genistein was obtained from ChemCon GmbH (Freiburg i. Br.), Lot No.CC-6700-26, as a
powder. DMSO was obtained from Sigma (St. Louis, MO), Lot No.80K3695 0.9% Sodium
Chloride for Injection, USP (saline) was manufactured by Abbott Laboratories (Lot No. 55-
5 199-DK). Sterile Water for Injection, USP (WFI) was manufactured by Lyphomed, Inc. (Lot
No. 390849).

Formulations: Test preparations (treatment groups) are summarized in Table 4.

For preparation of formulations 1- 4, HHT-NCI was weighed into vials and dissolved in
DMSO at the stated concentrations.

10 For formulation 5, the contents of a 10-mg vial of lyophilized CDDP (Cisplatin for Injection)
was resuspended with 10 mL WFI to produce a 1 mg/mL CDDP suspension.

For formulation 6, paclitaxel, prediluted in Cremaphor/EL and dehydrated alcohol to 6
mg/mL was further diluted to 3.3 mg/mL with WFI.

15 Formulations 7 and 8 were prepared by further diluting HHT-Clin to the stated
concentrations with WFI. Formulation 9 was undiluted HHT-Clin, used as received.

Formulation 10 was prepared by adding 1 mL of WFI to 100 mg of cytarabine as a
lyophilized powder.

Formulation 11 was prepared by adding DMSO to camptothecin at a concentration of 1
mg/mL.

20 Formulation 12 was made by adding 0.9% Sodium Chloride for Injection to a vial of 10 mg
of vinblastine lyophilized powder.

Formulations 13-17 were prepared by diluting the appropriate amount of each test agent into
saline (13- 2.5 mg/mL etoposide, 14- 7.5 mg/mL 5-fluorouracil, 15- 7.5 mg/mL amonafide,
16- 2.5 mg/mL colchicine, 17- 3.75 mg/mL 5-fluorouracil).

Formulation 18 was prepared by diluting 15 mg of genistein in 1 mL of DMSO.

Animals: Female C3H mice (Charles River Laboratories, Holister, CA), approximately 3 months old, were used for the study. The average body weight was approximately 25 g. Animals were maintained in isolator cages on a 12-hour light-and-dark cycle. Food and water were available *ad libitum*.

Tumors: The RIF-1 murine fibrosarcoma cell line was maintained in *in vitro* culture (Waymouth medium supplemented with 20% fetal bovine serum) at 37° C in a humidified 5 % CO₂ incubator. Log-phase RIF-1 cells were trypsinized and harvested from cell culture flasks to yield a concentration of 4 x 10⁶ cells/mL, then injected intradermally in a volume of 50 µL (equivalent to 2 x 10⁵ cells per injection) into both flanks of each mouse. Nine days later, when tumors reached approximately 100 mm³ in size, the animals were randomized to different treatment groups.

Treatment Groups: Treatment groups are summarized in Table 4. Four to five animals were assigned to each treatment group. The intraperitoneal injection volume was 100 µL. Intratumoral injections (50 µL) were made into one of the two tumors on each animal with the contralateral tumor serving as an untreated control. The oral administration volume was 100 µL. Combination treatments using two test agents were administered as two separate injections, with the second one following the first either immediately or after 30 minutes.

Evaluation of Tumor Volume Quadrupling Time: Tumors were measured three times weekly for up to 22 days with Vernier calipers. Tumor volume (cubic millimeters, mm³) was calculated according to the formula:

$$V = \pi / 6 \times D_1 \times D_2 \times D_3$$

in which D₁₋₃ are perpendicular diameters measured in millimeters (mm).

Tumor volume quadrupling time (TVQT), defined as the time required for a tumor to grow to four times its initial volume (at the time of treatment), was used as a study endpoint. The TVQT was determined for each treatment group and expressed in days as the

mean \pm standard error (SE).

Antitumor activity or modulation of tumor growth (as measured by delayed tumor growth, i.e. increases in TVQT values) by homoharringtonine administered as a single agent or in combination with other chemotherapeutics is presented in Table 5.

- 5 Results from eight separate experiments are included in this study. In experiment E010, tumors in untreated control animals quadrupled in size in an average of 7.2 days. Intraperitoneal administration of homoharringtonine from NCI at 5 mg/Kg had a TVQT of 14.5 days and intratumoral administration of homoharringtonine at that dose resulted in a TVQT of 15.6 days.
- 10 In experiment E011, untreated control animals quadrupled in size an average of 8.3 days while intraperitoneal administration of homoharringtonine from NCI at 2 mg/Kg extended the mean TVQT to 10.1 days, and the additional intraperitoneal administration of CDDP further extended the mean TVQT to 14.9 days. While paclitaxel (10 mg/Kg), alone, demonstrated a TVQT of 8.8 days, the addition of homoharringtonine (2 mg/kg) did not
- 15 change the TVQT, making paclitaxel the only agent with combinatorial activity less than that of homoharringtonine, alone.

Homoharringtonine from Hangzhou Minsheng Pharmaceutical Group (Hangzhou, China), formulated in sterile water at either 2 mg/Kg or 4 mg/Kg was used for the remainder of the combination studies.

- 20 At 2 mg/Kg, homoharringtonine had an average TVQT of 10.4 days in E026 while the untreated controls quadrupled in 7.4 days. Combination administration of cisplatin (4 mg/Kg) with homoharringtonine (2 mg/Kg) yielded a TVQT of 11.1 days, which was greater than homoharringtonine (TVQT= 10.4 days) or cisplatin (TVQT= 9.4 days), alone.

- In experiment E030, where untreated controls quadrupled in 6.7 days, homoharringtonine
- 25 treatment (2 mg/Kg) yielded a TVQT of 7.9 days and camptothecin or cytarabine gave

TVQT's of 9.4 or 7.6 days, respectively. Combination administration of homoharringtonine (2 mg/Kg) with camptothecin (6 mg/Kg) or cytarabine (400 mg/Kg) increased the TVQT's to 10.1 and 8.6 days, respectively.

In E032, where untreated controls quadrupled in 6.5 days, homoharringtonine at 4 mg/Kg had an average TVQT of 8.5 days. Administration of homoharringtonine (4 mg/Kg) in combination with 5-fluorouracil (30 mg/Kg) resulted in a TVQT of 17.9 days versus 13.6 days for 5-fluorouracil, alone. Combination administration of homoharringtonine (4 mg/Kg) and vinblastine (2 mg/Kg) yielded a TVQT of 10.9 days versus 8.6 days for vinblastine, alone. Combination administration of homoharringtonine (4 mg/Kg) and cisplatin (4 mg/Kg) or amonafide (30 mg/Kg) yielded TVQT's of 10.4 and 10.2 days, respectively, versus 9.9 and 7.6 days for those agents, alone. Homoharringtonine in combination with etoposide (10 mg/Kg) gave a TVQT of 8.7 days while etoposide, alone, was 8.5 days.

Orally administered colchicine (10 mg/Kg), in E033, yielded a TVQT of 6.3 days, while untreated controls and homoharringtonine (4 mg/Kg) gave TVQT's of 7.8 and 8.3 days. Homoharringtonine in combination with colchicine at these doses increased the TVQT to 9.4 days.

In E036, genistein (60 mg/Kg) in combination with homoharringtonine (4 mg/Kg) had a TVQT of 9.2 days, which was greater than that of genistein, alone (7.1 days).

There were animal deaths in some groups that were recorded as follows. Three of four mice died after treatment of homoharringtonine obtained from NCI and formulated in DMSO at 1.25 mg/mL. Two of five mice died after receiving this formulation intratumorally. Four of four mice died after treatment of this same homoharringtonine formulated to 2.5 mg/mL in DMSO. The combination of homoharringtonine (0.5 mg/mL) in DMSO with paclitaxel (2.5 mg/mL) was lethal to two of four mice, and the combination of homoharringtonine (0.5 mg/mL) in DMSO with cisplatin (1 mg/mL) was lethal to one of four mice. The combination of homoharringtonine (1 mg/mL) in vinblastine (0.5 mg/mL) was lethal to one of four mice given that treatment, and the combination of homoharringtonine (1 mg/mL) and

genistein (15 mg/mL) was lethal to two of five mice.

In summary, intraperitoneal administration of homoharringtonine had antitumor activity, i.e. modulated tumor growth, in the RIF-1 murine fibrosarcoma tumor model. Intraperitoneal administration of homoharringtonine in combination with cisplatin, cytarabine, camptothecin, vinblastine, etoposide, 5-fluorouracil, amonafide, colchicine and genistein had antitumor activity levels greater than homoharringtonine alone, or the individual test agents. The best combinatorial activity used 5-fluorouracil, amonafide and vinblastine. Homoharringtonine in combination with paclitaxel had antitumor activity less than homoharringtonine alone. Homoharringtonine obtained from NCI and formulated in DMSO showed some lethal toxicity while homoharringtonine obtained from Hangzhou Minsheng Pharmaceutical Group (Hangzhou, China) and formulated in sterile water for use in humans did not show lethal toxicity at the doses used.

Table 4
Summary of Treatment Groups

Formulation	Treatment	Concentration (mg/mL)	Route of Administration	Injection Volume (μL)
1	HHT-NCI in DMSO	1.25	IP	100
2	HHT-NCI in DMSO	2.5	IP	100
3	HHT-NCI in DMSO	2.5	IT	50
4	HHT-NCI in DMSO	0.5	IP	100
5	CDDP in WFI	1	IP	100
6	Paclitaxel in WFI	2.5	IP	100
7	HHT-Clin in WFI	0.5	IP	100
8	HHT-Clin in WFI	0.25	IP	100

9	HHT-Clin in WFI	1	IP	100
10	Cytarabine in WFI	100	IP	100
11	Camptothecin in DMSO	2.5	IP	100
12	Vinblastine in saline	0.5	IP	100
13	Etoposide in saline	2.5	IP	100
14	5-Fluorouracil in saline	7.5	IP	100
15	Amonafide in saline	7.5	IP	100
16	Colchicine in saline	2.5	PO	100
17	5-Fluorouracil in saline	3.75	IP	100
18	Genistein in DMSO	15	IP	100

Table 5

Effect of Homoharringtonine and Homoharringtonine in Combination with Other Chemotherapeutics on RIF-1 Tumor Growth in C3H Mice

Exp. #	Formulation	Treatment	# of Tumors	TVQT (days) (Mean \pm SE)
E010	-	Untreated control	8	7.2 \pm 0.1
E010	1	HHT-NCI (5 mg/Kg)	2*	14.5 \pm 0.9
E010	2	HHT-NCI (10 mg/Kg)	0*	All Died
E010	3	HHT-NCI (5 mg/Kg)	3*	15.6 \pm 1.8
E011		Untreated control	8	8.3 \pm 0.4
E011	4	HHT-NCI (2 mg/Kg)	8	10.1 \pm 0.4

E011	5	CDDP (4 mg/Kg)	8	12.9 ± 1.1
E011	4,5	HHT-NCI -30'- CDDP	6*	14.9 ± 0.8
E011	6	Paclitaxel (10 mg/Kg)	8	8.8 ± 0.4
E011	4,6	HHT -30'- Paclitaxel	4*	8.8 ± 0.4
E026	-	Untreated control	8	7.4 ± 0.3
E026	7	HHT-Clin (2 mg/Kg)	8	10.4 ± 1.0
E026	5	CDDP (4 mg/Kg)	8	9.4 ± 0.5
E026	7,5	HHT-Clin + CDDP	8	11.1 ± 0.4
E026	7,5	HHT-Clin -30'- CDDP	8	10.1 ± 0.4
E028	-	Untreated control	8	8.7 ± 0.5
E028	8	HHT-Clin (1 mg/Kg)	8	9.2 ± 0.7
E028	9	HHT-Clin (4 mg/Kg)	8	10.1 ± 0.4
E030	-	Untreated control	8	6.7 ± 0.4
E030	7	HHT-Clin (2 mg/Kg)	8	7.9 ± 0.3
E030	10	Cytarabine (400 mg/Kg)	8	7.6 ± 0.2
E030	7,10	HHT-Clin + Cytarabine	8	8.6 ± 0.4
E030	11	Camptothecin (6 mg/Kg)	8	9.4 ± 0.4
E030	7,11	HHT-Clin + Camptothecin	8	10.1 ± 0.6
E032	-	Untreated control	8	6.5 ± 0.6
E032	9	HHT-Clin (4 mg/Kg)	8	8.5 ± 0.5
E032	5	CDDP (4 mg/Kg)	8	9.9 ± 0.6
E032	9,5	HHT-Clin + CDDP	8	10.4 ± 0.4




E032	12	Vinblastine (2 mg/Kg)	8	8.6 ± 0.4
E032	9,12	HHT-Clin + Vinblastine	6*	10.9 ± 0.4
E032	13	Etoposide (10 mg/Kg)	8	8.5 ± 1.0
E032	9,13	HHT-Clin + Etoposide	8	8.7 ± 0.5
E032	14	5-Fluorouracil (30 mg/Kg)	8	13.6 ± 1.9
E032	9,14	HHT-Clin + 5-Fluorouracil	8	17.9 ± 0.7
E032	15	Amonafide (30 mg/Kg)	8	7.6 ± 0.4
E032	9,15	HHT-Clin + Amonafide	8	10.2 ± 0.5
E033	-	Untreated control	8	7.8 ± 0.6
E033	9	HHT-Clin (4 mg/Kg)	8	8.3 ± 0.4
E033	16	Colchicine (10 mg/Kg)	8	6.3 ± 0.3
E033	9,16	HHT-Clin + Colchicine	8	9.4 ± 0.5
E033	17	5-Fluorouracil (15 mg/Kg)	8	6.7 ± 0.4
E033	9,17	HHT-Clin + 5 Fluorouracil	8	8.6 ± 0.3
E036	-	Untreated control	8	6.8 ± 0.4
E036	18	Genistein (60 mg/Kg)	8	7.1 ± 0.4
E036	9,18	HHT-Clin + Genistein	6*	9.2 ± 0.5

* Animal deaths occurred in these groups. See text for details.

WE CLAIM:

1. A method of treatment of a host with a cellular proliferative disease, comprising contacting said host with a cephalotaxine and an antiproliferative agent, each in an amount sufficient to modulate said cellular proliferative disease.
- 5 2. The method according to claim 1, wherein said cephalotaxine comprises the butanediocate ester cephalotaxine, 4-methyl-2-hydroxy-2-(4-hydroxy-4- methyl pentyl).
3. The method according to claim 1, wherein said cephalotaxine comprises a homoharringtonine analog.
- 10 4. The method according to claim 1 wherein said antiproliferative agent comprises an agent that interacts with nucleic acids.
- 5 5. The method according to claim 1 wherein said antiproliferative agent comprises an alkylating agent, an intercalating agent, a metal coordination complex, a pyrimidine nucleoside, a purine nucleoside, an inhibitor of nucleic acid associated enzymes, or an inhibitor of nucleic acid associated proteins.
- 15 6. The method according to claim 1 wherein said antiproliferative comprises cisplatin, cytarabine, camptothecin, vinblastine, etoposide, 5-fluorouracil, amonafide, colchicine, or genistein.
7. A method according to claim 1 wherein said cephalotaxine is administered before the administration of said antiproliferative agent.
- 20 8. A method according to claim 1 wherein said cephalotaxine is administered during the administration of said antiproliferative agent.

9. A method according to claim 1 wherein said cephalotaxine is administered after the administration of said antiproliferative agent.
10. The method of claim 1 wherein the modulation on said disease with said composition is greater than that for said antiproliferative agent alone.
- 5 11. A composition comprising a cephalotaxine and an antiproliferative agent.
12. The composition of claim 11 wherein said cephalotaxine comprises homoharringtonine.
13. The composition of claim 11 wherein said antiproliferative agent comprises cisplatin, cytarabine, camptothecin, vinblastine, etoposide, 5-fluorouracil, amonafide, colchicine, or genistein.
- 10 14. Use of a cephalotaxine and an antiproliferative agent in the formulation of a medicament for the treatment of a cellular proliferative disease.

**FIGURE 1**




FIGURE 2




FIGURE 3

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68098 A3

(51) International Patent Classification?: **A61K 31/55.**
A61P 35/00

(21) International Application Number: PCT/US01/08480

(22) International Filing Date: 15 March 2001 (15.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/189,699 15 March 2000 (15.03.2000) US

(71) Applicant (for all designated States except US): **CHEM-GENEX THERAPEUTICS, INC.** [US/US]: 34700 Campus Drive, Fremont, CA 94555 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BROWN, Dennis, M.** [US/US]: 100 San Mateo Drive, Menlo Park, CA 94025 (US). **MICHAELS, Shawnya, J.** [US/US]: 206 Dolores Street, El Granada, CA 94018 (US).

(74) Agents: **TRECARTIN, Richard, F.** et al.; Flehr Hohbach Test Albritton & Herbert LLP, 4 Embarcadero Center, Suite 3400, San Francisco, CA 94111-4187 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
6 June 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CEPHALOTAXINE ALKALOID COMBINATION COMPOSITIONS AND USES THEREOF

(57) Abstract: A method of treatment of a host with a cellular proliferative disease, comprising contacting the host with a cephalotaxine and an antiproliferative agent, each in an amount sufficient to modulate said cellular proliferative disease, is described. In some embodiments, the cephalotaxine comprises homoharringtonine (cephalotaxine, 4-methyl-2-hydroxy-2-(4-hydroxy-4-methyl pentyl) butanedioclate ester). Antiproliferative agents of the invention comprise alkylating agents, intercalating agents, metal coordination complexes, pyrimidine nucleosides, purine nucleosides, inhibitors of nucleic acid associated enzymes and proteins, and agents affecting structural proteins and cytoplasmic enzymes.

WO 01/68098 A3



INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/08480

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K31/55 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EMBASE, EPO-Internal, CHEM ABS Data, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document; with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	C. VISANI ET AL: "Effects of homoharringtonine alone and in combination with alpha interferon and cytosine arabinoside on "in vitro" growth and induction of apoptosis in chronic myeloid leukemia and normal hematopoietic progenitors" LEUKEMIA, vol. 11, 1997, pages 624-628, XP008000008 abstract --- -/--	1,3,6, 10-14

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

7 March 2002

Date of mailing of the international search report

22/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Siatou, E

INTERNATIONAL SEARCH REPORT

Int. .tional Application No

PCT/US 01/08480

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages.	Relevant to claim No.
X	<p>DATABASE EMBASE 'Online! ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL; YUZHU Z. ET AL: "Homoharringtonine, cytarabine and aclarubicin (HAA) combination chemotherapy for acute myeloid leukemia (AML)." retrieved from STN Database accession no. 1998384948 XP002192356 abstract & CHINESE JOURNAL OF CLINICAL ONCOLOGY, (1998) 25/10 (758-759). ,</p>	1,3,6, 10-14
X	<p>----- DATABASE EMBASE 'Online! ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL; LASTER JR. W.R. ET AL: "Therapeutic synergism (TS) of homoharringtonine (H) plus 5-fluorouracil (FU) against leukemia P388 (P388/O) and ARA-C-resistant P388 (P388/ARA-C)." retrieved from STN Database accession no. 82182588 XP002192357 abstract & PROCEEDINGS OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, (1982) VOL. 23/- (NO. 786). CODEN: PAACA3,</p>	1,3,6, 10-14
X	<p>----- ZHANG ET AL: "Inhibitory effects of homoharringtonine and hydroxycamptothecin in combination with other agents on cancer cell growth" ASIA PACIFIC JOURNAL OF PHARMACOLOGY, SINGAPORE UNIVERSITY PRESS, SG, vol. 7, 1992, pages 191-195, XP002112006 ISSN: 0217-9687 page 194, right-hand column abstract</p>	1,3,6, 10-14
A	<p>----- TAKANO I ET AL: "Ester-type cephalotaxus alkaloids from cephalotaxus harringtonia var. drupacea" PHYTOCHEMISTRY, PERGAMON PRESS, GB, vol. 44, no. 4, February 1997 (1997-02), pages 735-738, XP004292875 ISSN: 0031-9422 the whole document</p> <p>----- -/--</p>	1-14

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/08480

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TAKANO I ET AL: "NEW OXYGENATED CEPHALOTAXUS ALKALOIDS FROM CEPHALOTAXUS HARRINGTONIA VAR. DRUPACEA" JOURNAL OF NATURAL PRODUCTS, XX, XX, vol. 59, no. 12, December 1996 (1996-12), pages 1192-1195, XP000990654 ISSN: 0163-3864 abstract</p>	1-14